



DementiaUK
Helping families face dementia

Recognising the later stages of dementia and moving towards end of life care



Dementia is a progressive neurological condition that is life-limiting. This means that everyone with dementia will die with or from it, although many people die as a result of other conditions (such as cancer or heart failure) before they reach its later stages.

Whether a person has dementia alone or other health conditions alongside, it is important that plans are made to ensure they receive good care throughout their life. Towards the end of life, this may include a shift away from treatments that may cause distress or have limited benefit, and more towards a focus on comfort.

The end of life stage can be hard to identify and accept – by the person with dementia, if they are aware of what is happening, and their family members, who often play a vital role in supporting them. However, good end of life care can enhance the person's quality of life and ensure their needs are met; this can also give their family and friends peace of mind that they are being cared for well.

Amongst professionals, it is usually understood that a person will move towards end of life care when it is thought that they are within the last months or years of life. However, this is very individual and is often difficult to predict, so it is better to focus on their needs rather than on how long they may have left to live.

Identifying the later stages of dementia and the end of life

Signs that a person with dementia might be approaching the end of life include:

- significantly reduced appetite: loss of interest in food and drink, or refusing it completely
- becoming increasingly frail
- weight loss



- swallowing difficulties (for example, coughing/throat-clearing during or after eating or drinking; taking a long time to chew food; choking on food; or storing food in their mouth)
- frequent infections
- reduced mobility, perhaps requiring care in a bed or chair
- frequent falls
- reduced communication
- becoming more withdrawn, or agitated and/or restless
- sleeping more, seeming drowsy and less aware
- difficulties controlling their bladder and/or bowel (incontinence)
- needing help with most daily activities

If the person you care for is showing a number of these signs or you have any concerns, make an appointment with their GP – there may be an underlying cause that could be treated or managed better, such

4 Dementia UK

as pain or delirium. Please see Sources of support on p17-18 for our leaflets on these.

Decision-making towards the end of life

If the person you care for is entering the later stages of dementia or another condition, it is important to think about how their end of life care should be provided. Making a clear, documented plan will help everyone work together to support the person in the later stages of their life. It can bring a sense of calm and control to emotional situations, and avoid having to make decisions in a crisis, which may be distressing.

If possible, the person with dementia should be involved in making decisions about their future care before they reach a stage where they are unable to do so. This may include making:

- an advance care plan: this covers how they would like to be cared for and where they would like to die, as well as information about subjects like their Will and funeral. You can download a template at [▶ dementiauk.org/wp-content/uploads/dementia-uk-advance-care-plan-template.pdf](https://dementiauk.org/wp-content/uploads/dementia-uk-advance-care-plan-template.pdf)
- an advance decision to refuse treatment (ADRT): also known as an ‘advance decision’ or ‘living Will’, this states which treatments they do not wish to receive, such as antibiotics, ventilation and/or resuscitation (CPR). You can find out more at [▶ nhs.uk/conditions/end-of-life-care/advance-decision-to-refuse-treatment/](https://nhs.uk/conditions/end-of-life-care/advance-decision-to-refuse-treatment/)

You might want to ask the GP for help with putting together an advance care plan or ADRT. They may be able to advise on local services that could support you. You can also contact our dementia specialist Admiral Nurses for advice – see p16 for more information.

If the person with dementia is reaching the end of life and has not previously made their wishes and preferences known, it should not be assumed that they are now unable to do so. However, in the later stages of dementia, they may lack the mental capacity to make some decisions. In this case, other people may need to make these decisions on their behalf, in their best interests. For our information on mental capacity and decision-making, please see Sources of support on p18.

Some of the decisions you may want to think about include:

Resuscitation

Making the decision that someone should not be resuscitated if their heart stops can be extremely difficult. But resuscitation is often unsuccessful for a person in the advanced stages of dementia and could cause distress and trauma. If it is unsuccessful, it may also lead to them dying in an unfamiliar place, such as an ambulance or A&E department, rather than in their own home or care home.

For this reason, the person with dementia (if they still have mental capacity), their health and/or social care professionals and you as their next of kin and/or carer may decide in advance that it is in their best interests not to attempt resuscitation at this stage.

The NHS has more information to help you decide whether resuscitation is appropriate for the person you care for as they near the end of life: [▶ nhs.uk/conditions/do-not-attempt-cardiopulmonary-resuscitation-dnacpr-decisions](https://www.nhs.uk/conditions/do-not-attempt-cardiopulmonary-resuscitation-dnacpr-decisions)

Eating and drinking

At the end of life, people may have a reduced appetite or not want to eat or drink at all. This can be hard to accept, but it is a natural part of the dying process. Invasive interventions like feeding through a tube

6 Dementia UK

or drip are unlikely to extend the person's life or improve their quality of life, and may cause distress.

Instead, you and the person's care team might decide to focus on careful hand-feeding and giving food and drink for comfort and enjoyment, rather than nutrition. For example, you could:

- offer food and fluids from a teaspoon – carefully and in small amounts
- give foods with an appealing taste and texture
- give just a few mouthfuls at a time
- use ice chips or artificial saliva spray to keep their mouth comfortable
- ask the person's GP for a referral to a speech and language therapist for advice, especially if they have difficulty swallowing

Please see Sources of support on p17-18 for details of our leaflets on eating and drinking; hydration for a person with dementia; and understanding dying, which may be helpful at this stage.

Infections

Frequent infections are common in the end stages of life, especially chest infections, pneumonia and urinary tract infections (UTIs). It is important to consult the person's GP if they are showing signs of infection to discuss the best course of action.

If the person with dementia can still swallow, they may be prescribed liquid antibiotics to be taken by mouth. Even if they do not cure the infection, they could help to relieve symptoms and reduce discomfort and distress. However, the GP may feel that antibiotics are not appropriate – for example, if the person has swallowing difficulties and antibiotics would have to be administered by drip



– and prescribe other types of medication to help keep them as comfortable and peaceful as possible.

Frailty, mobility and falls

In the later stages of dementia, many people become increasingly frail and experience mobility problems and falls. This is often part of the progression of the condition (or of other conditions that they have alongside dementia).

If the person you care for is having falls, it is a good idea to speak to their GP in case there is another cause that could be treated or managed, such as low blood pressure, problems with their balance, or poor eyesight.

You and the person's GP and/or other health and social care professionals may want to discuss whether it is right for them to be admitted to hospital if they fall. For example, if the person may have a fracture or other serious injury, they will require hospital assessment and/or treatment. Otherwise, their health professional may feel they can be supported to stay at home or in their care home.

Becoming frail and less mobile may lead to other health problems – for instance, if the person is spending long periods of time in a bed or chair, they may be more likely to develop constipation and pressure sores. The person’s GP may be able to refer them to a district nurse for advice and support on keeping them comfortable.

Where the person will be cared for at the end of life

It is important to think about where the person with dementia will be cared for at the end of life and any wishes they have expressed about where they would like to die. It is also important to take your own feelings into account – for example, whether you think you can continue to care for them at home or would find the memory of them dying at home traumatic.

People at the end of life often need 24-hour care. It may be extremely difficult to provide this care yourself, so if you and the person with dementia feel it is important for them to remain at home, you may want to consider support from professional carers.

Social Services can arrange a needs assessment for the person with dementia and a carer’s assessment for the family carer to explore the support options available, such as respite care, adaptations to make the home safer and more comfortable, paid home carers, or home-based palliative care. For information on applying for these assessments, please see Sources of support on p18-19.

As the person’s care needs increase, you may want to consider a move to a care home or nursing home. This can be a very big and emotional decision, and you may feel you are letting the person down, but care homes can provide the extra support they need so you can focus on spending time together. Our leaflet on considering a care home may be helpful – please see Sources of support on p18.

If there is a hospice in your area, they may be able to support you in caring for the person with dementia at the end of their life. This might include advice on symptom management and supportive care that involves family members and helps the person stay as comfortable as possible in a familiar and reassuring environment.

While some people are admitted to a hospice at the end of life, it is more likely that the hospice nurses will support the person to remain at home, unless they have acute and intense symptoms that cannot be managed at home – which may be the result of another condition like cancer. Your GP can advise on any local hospices and palliative care provision that may be able to support you.

Keep in mind that decisions about the person's care may need to be revisited and adapted as circumstances change – for example, it may be necessary for them to be admitted to hospital – but even if the person is being cared for elsewhere, you can still play a vital part in helping to meet their needs.

Emotional care

Although a person's physical abilities, memory and ability to communicate often deteriorate as their dementia advances, it is thought that feelings remain intact. The person may not be able to tell you how they are feeling – although their behaviour may be a good indicator – but it is important that they continue to feel loved, safe and secure.

The senses can be a powerful way to maintain connections and give comfort and pleasure. This could include:

- using scents that they find appealing or that have been favourites in the past, such as a perfume, aftershave or soap that they like to use

- looking at photos together
- playing music that they enjoy listening to
- giving them a gentle hand massage
- compiling a memory box to look through and handle together
- brushing their hair
- giving them small amounts of their favourite food and drink
- giving them something comforting to hold, such as a fluffy blanket or soft toy
- talking about happy occasions in their past

Compiling a life story

If the person with dementia has not already created a life story, you may want to do this before or as they enter the later stages of dementia. This is a record of things that are important to them, such as their family and friends; work history; important occasions in their past; likes and dislikes; religious, cultural and spiritual beliefs; and future wishes.

A life story can be recorded in a book, a computer document, an app, a memory box, a collage and/or a video recording. You can make it together with the person with dementia or, if their condition is too advanced, on their behalf. It can then be shared with professionals and carers to help them connect with the person in an individual and meaningful way. For more information and our life story template, please see Sources of support on p17.

Faith, spirituality and culture

For many people, faith, spirituality and culture play a big part in their life. It is important that these values and needs are



considered at the end of life, such as through religious practices or ceremonies, traditional rituals, songs/music, meditation, or clothing. For example:

- If the person with dementia follows a religion, a visit, blessing or prayer from a religious leader or hospital/hospice chaplain may provide comfort and reassurance
- If their religious or cultural community has rituals around the end of life, such as what should happen immediately before or after they die (eg receiving the last rites) they may need assurance that these will be followed
- There may be particular people that the person wants to see before they die to ensure they feel at peace

The best way to find out what is important to the person with dementia is to ask them, if possible; if not, their family, other people who are close to them, or a religious or cultural leader may be able to help. You could include this information in their care plan or life story (see p16-17).

Young onset dementia and the end of life

Most of the advice around end of life care with dementia also applies to young onset dementia (where symptoms develop before the age of 65), and important issues like advance care planning and resuscitation should still be considered, even though the person may appear in better physical health and less frail than an older person.

However, there are likely to be additional considerations linked to the person's age. Depending on their level of understanding, the person may have worries that are specific to their life stage – for example, they may be distressed about leaving their children or anxious about their partner having to manage financial commitments like a mortgage alone.

The emotional impact of the person dying at a younger age, the loss of the future they had imagined, and the prospect of having to manage without them can be very difficult for families. Losing the person's salary or benefits can be another source of considerable stress for family members.

You may also need to consider issues like:

- the ability of their partner and other family members to take time off work as the person with dementia nears the end of life
- the person's wishes for the end of life if they are known, eg where they would prefer to die
- whether it is appropriate for the person to stay at home to die if there are children or young people living there
- if the person does not remain at home, where they will be cared for towards the end of life (some care homes do not accept residents under the age of 65; others may have difficulty meeting the needs of a younger person)

- what will happen if the person is admitted to hospital, eg whether they would be admitted to a general ward or an older persons' ward
- whether there is hospice care to meet the person's needs and those of their family – either to offer end of life support at home, or to admit the person to the hospice as a home-from-home that may be less intimidating for them and their family
- whether their children should be able to visit the person at the end of life
- who will look after their children if you decide that they should not be present at the end of life

If you work, it is a good idea to make your employer aware that you are caring for someone who is at the end of life so you can access carer's leave to ease any worries about taking time off work. Many employers continue to pay the employee while they are taking carer's leave, although they do not have to. You may also be able to access bereavement support through your employer.

Time together as a whole family, including the person who is dying if this is possible and reflects their needs and wishes, can be important in making lasting memories and ensuring everyone feels involved and valued. Working together on a life story (see p17) can help maintain connections between parent and child, as well as other family members.

A child or young person facing the imminent death of a parent is likely to need a lot of support. Conversations about the end of life can be very difficult, and you may worry about how much the child will understand and how they will react. But speaking honestly can help them to feel supported.

14 Dementia UK

These tips may help you support a child who is facing the loss of a parent:

- Give clear, simple and accurate information that is appropriate to the child's age and level of understanding
- Provide information in small chunks to allow the child time to process and understand it
- Ask if they have any questions and respond with the information they need at the time – you can expand on it later
- Keep to familiar routines (eg after-school clubs, mealtimes, bedtime) to help the child feel secure and maintain a sense of normality
- Let them help with caring tasks if they want to – even small things like making the bed or, for an older child, preparing a meal, could help them feel more involved
- Involve them in making decisions to give them some control in a world that may feel out of control – this could be as simple as helping choose what their parent will wear
- Provide the child with plenty of reassurance that they are loved and there are people who will be there for them
- Help the child make memories, for example by making memory boxes, photo books or drawings, and taking photos and videos with the person with dementia
- Talk to the child's school or college about what is happening so they understand that they may need extra support or time off at short notice



- The child may benefit from pre- and post-bereavement talking therapy and access to support services – you can ask their school or GP what is available

It is important that the person with young onset dementia and their family have professional support as they think about the later stages of dementia and make decisions about their wishes for end of life care. Please see Sources of support on p16-19 for information about people and organisations that could help.

Sources of support

As a carer or family member of a person at the end of life, it is vital that you feel well supported and able to look after your own needs. You may wish to seek help from:

- other family members or friends
- carers' support groups (in person or online)
- a local hospice
- health or social care professionals, eg your GP or a counsellor
- religious or community leaders
- your child's school

You can also contact our free Helpline to speak to a dementia specialist Admiral Nurse: please call **0800 888 6678** (Monday-Friday 9am-9pm, Saturday and Sunday 9am-5pm, every day except 25th December), or email [▶ helpline@dementiauk.org](mailto:helpline@dementiauk.org)

If you prefer, you can book a phone or video appointment with an Admiral Nurse at a time to suit you: visit [▶ dementiauk.org/book](https://dementiauk.org/book)

Dementia UK resources

Advance care planning

[▶ dementiauk.org/advance-care-planning](https://dementiauk.org/advance-care-planning)

Advance care plan template

[▶ dementiauk.org/care-plan-template](https://dementiauk.org/care-plan-template)

A stay in hospital

[▶ dementiauk.org/changes-in-care-a-stay-in-hospital](https://dementiauk.org/changes-in-care-a-stay-in-hospital)

The carer's assessment

➤ dementiauk.org/the-carers-assessment

Continence

➤ dementiauk.org/continence

Delirium

➤ dementiauk.org/delirium

Eating and drinking

➤ dementiauk.org/eating-and-drinking

Falls

➤ dementiauk.org/dementia-and-falls

Frailty and dementia

➤ dementiauk.org/frailty

Good hydration for a person with dementia

➤ dementiauk.org/hydration

Grief, bereavement and loss

➤ dementiauk.org/bereavement

Lasting power of attorney

➤ dementiauk.org/lasting-power-of-attorney

Life story work

➤ dementiauk.org/life-story

Life story template

➤ dementiauk.org/life-story-template

Looking after yourself as a carer

➤ dementiauk.org/looking-after-yourself

Mental capacity and decision-making

➤ dementiauk.org/capacity-decision-making

Moving into a care home

➤ dementiauk.org/changes-in-care/choosing-a-care-home

Pain and dementia

➤ dementiauk.org/pain

Supporting children and adolescents when a parent has young onset dementia

➤ dementiauk.org/supporting-children-and-adolescents

Understanding dying

➤ dementiauk.org/understanding-dying

Young onset dementia: choosing a care home

➤ dementiauk.org/young-onset-dementia-choosing-a-care-home

Other resources

Carer's assessment – find your local social services

➤ nhs.uk/service-search/other-services/Local-Authority-Adult-Social-Care/LocationSearch/1918

Carer's leave

➤ gov.uk/carers-leave

Child Bereavement UK

➤ childbereavementuk.org

Helpline: 0800 028 8840

Compassion in Dying: living Will advice

➤ compassionindying.org.uk



Dying Matters

➤ dyingmatters.org

Hospice UK

➤ hospiceuk.org

Marie Curie: care and support through terminal illness

➤ mariecurie.org.uk

Helpline: 0800 090 2309

The needs assessment

➤ gov.uk/apply-needs-assessment-social-services

Resuscitation Council UK: ReSPECT plan for emergency care and CPR

➤ resus.org.uk/respect/respect-patients-and-carers

**To speak to a dementia specialist Admiral Nurse
about any aspect of dementia:**

Contact our Helpline:
0800 888 6678 or [📧 helpline@dementiauk.org](mailto:helpline@dementiauk.org)

Book a virtual appointment:
[📅 dementiauk.org/book](https://dementiauk.org/book)

**Our charity relies entirely on donations to fund our
life-changing work. If you would like to donate to help us
support more families:**

- Call **0300 365 5500**
- Visit [📍 dementiauk.org/donate](https://dementiauk.org/donate)
- Scan the QR code

Thank you.



DementiaUK
Helping families face dementia



dementiauk.org • info@dementiauk.org

Publication date: May 2024. Review date: May 2026. © Dementia UK 2024

Dementia UK, 7th Floor, One Aldgate, London EC3N 1RE
Dementia UK is a registered charity in England and Wales (1039404) and Scotland (SC 047429).